

DETERMINATION OF PARTICULATE RESUSPENSION AND MIXING NEAR AN OCEAN OUTFALL USING
HYPERSPECTRAL OCEAN COLOR IMAGERY

Michael Hamilton, Curtiss O. Davis

Jet propulsion Laboratory
California Institute of Technology
4800 Oak Grove Drive
Pasadena, CA 91109
(818) 354 58G9
(81 8) 393-6720 (fax)

In-water apparent optical properties, spectral reflectance at the **sea** surface, and an **AVIRIS** image were acquired **simultaneously** in the vicinity of an ocean outfall. Sampling was subsequent to several days of storms and nearby mooring data suggest a resuspension of outfall sediments in response to increased currents. **The** sediment plume is clearly visible in a **derivative** image, centered at the **shoulder** of the sediment reflectance feature at 590 **nm**.

Instability waves **caused** by the relative motion of two fluids of different density are **seen** along the interface **between** water containing resuspended particles and the clearer offshore water mass. The rate of mixing of **the** particulates **into** the surrounding waters can be estimated from the spatial scale of **interfingering** along the front.

This result suggests that coastal monitoring with **high-spectral** and spatial resolution optical devices may be **useful** in determining the mixing and **areal extent** of sewage spills and resuspended **effluent** and the **threat** posed to nearby beach **areas**.